

# JUSTIN A. SAUNDERS, M.D.

DOCTORS PARK, SUITE 101  
1517 NICHOLASVILLE ROAD  
LEXINGTON, KENTUCKY 40503  
TEL (859) 277-4403  
FAX (859) 277-4405

JOSEPH H. SAUNDERS, M.D.  
(DECEASED)

JUSTIN A. SAUNDERS, M.D.  
PLASTIC AND RECONSTRUCTIVE SURGERY  
OF EYELIDS, TEAR DUCTS AND ORBIT  
justinsaundersmd@gmail.com

JOHN H. SAUNDERS, M.D.  
(RETIRED)

## Consent Form for Incision and Drainage of Chalazion

### *Condition and Proposed Treatment*

Your ophthalmologist has evaluated you and diagnosed you with a chalazion, which is a localized inflammatory response involving sebaceous glands of the eyelid that occurs when the gland duct is obstructed. A chalazion may resolve spontaneously or with warm compresses, lid scrubs, and lid massage. When there is no improvement, the chalazion may be incised and drained. After local anesthesia, a chalazion instrument is put in place and an incision is made in the inner aspect of the eyelid. The contents of the chalazion are then carefully drained with a curette followed by gentle pressure or heat to control any bleeding.

### *Alternatives to Surgery*

1. Lid Hygiene – Warm compresses, lid massage and scrubs; may not improve chalazion if deep.
2. Steroid Injection – May require more than one injection. Can result in depigmentation of the eyelid, steroid deposits at the injection site, or in rare instances occlusion of retinal and choroidal blood vessels with possible loss of vision
3. No Treatment – I may choose no treatment and tolerate the chalazion.

### *Risks and Complications*

No procedure is entirely risk free. Adverse effects from incision and drainage of chalazion may include:

1. Infection – Infections can be treated with topical or oral antibiotics
2. Bleeding – Normally controlled with gentle pressure or heat cautery at the incision site.
3. Pain – Minimal and resolves with healing of incision.
4. Recurrence – Chalazion may recur if incomplete excision.
5. Loss of lashes in the involved area
6. Eyelid notching in the area of the inflammation
7. Damage to the globe from the scalpel, needle used to inject the anesthetic, or cautery instrument.
8. Vision loss, including blindness.

*Consent for Treatment*

By signing below I acknowledge that I have read and understand the above, and have had my questions answered by the surgeon to my satisfaction. I consent to the incision and drainage of the chalazion on the \_\_\_\_\_ (state "upper" or "lower") lid of my \_\_\_\_\_ (state "right" or "left") eye.

\_\_\_\_\_  
Patient Name

\_\_\_\_\_  
Patient Signature

\_\_\_\_\_  
Date